

Ground State of Supermembrane on PP-wave

Noriko Nakayama^{†a}, Katsuyuki Sugiyama^{*} and Kentaroh Yoshida^{†b}

[†]*Graduate School of Human and Environmental Studies,
Kyoto University, Kyoto 606-8501, Japan.*

E-mail: ^anakayama@phys.h.kyoto-u.ac.jp

^byoshida@phys.h.kyoto-u.ac.jp

^{*}*Department of Fundamental Sciences,
Faculty of Integrated Human Studies,*

Kyoto University, Kyoto, 606-8501, Japan.

E-mail: sugiyama@phys.h.kyoto-u.ac.jp

Abstract

We consider the ground state of supermembrane on the maximally supersymmetric pp-wave background by using the quantum-mechanical procedure of de Wit-Hoppe-Nicolai. In the pp-wave case the ground state has non-trivial structure even in the zero-mode Hamiltonian, which is identical with that of superparticles on the pp-wave and resembles supersymmetric harmonic oscillators. The supergravity multiplet in the flat case comes to split with a certain energy difference. We explicitly construct the unique supersymmetric ground-state wave function of the zero-mode Hamiltonian, which is obviously normalizable. Moreover, we discuss the nonzero-mode Hamiltonian and construct an example for the ground-state wave function with a truncation of the variables. This special solution seems non-normalizable but its L^2 -norm can be represented by an asymptotic series.

Keywords: supermembrane, matrix theory, M-theory, pp-wave

Contents

1	Introduction	1
2	Supercharges and Superalgebra	3
2.1	Setup	3
2.2	$Spin(7) \times U(1)$ Formalism	6
3	Supersymmetric Quantum Mechanics of Zero-mode Hamiltonian	12
4	Nonzero-mode Ground-state Wave Function	19
5	Conclusions and Discussions	22
A	Proof for the $SO(3) \times SO(6)$ Invariance of the Ground State	24

1 Introduction

Recently, strings and M-theory on the pp-wave have been greatly focused on. The maximally supersymmetric pp-wave background in eleven dimensions [1] (KG solution) is one of the candidates for M-theory background. This background is achieved from the geometry of $AdS_4 \times S^7$ or $AdS_7 \times S^4$ [2] via the Penrose limit [3]. Similarly, the maximally supersymmetric type IIB pp-wave background was found [4], on which the string theory is exactly solvable [5,6,7]. Moreover, this string theory was used in the context of the AdS/CFT correspondence between the string spectra and the operators in Yang-Mills side [8].

Furthermore, the matrix model on the pp-wave was proposed in Ref. [8]. Originally, its action is obtained from the study of superparticles. In the flat space, the supermembrane theory [9,10,11] leads to the matrix model through the matrix regularization [11], which is considered as a candidate for M-theory [12]. Following the same procedure, the matrix model on the pp-wave can be also obtained from the supermembrane [13]. It was also shown that the superalgebra of the matrix model agrees with that of the supermembrane [14] as in the flat case [15].

In our previous works, we studied the supermembrane on the pp-wave from several aspects. In Ref. [16] we investigated BPS conditions of the supermembrane. The BPS states in the matrix model on the pp-wave have been intensively studied [17]. It was also shown in Ref. [8]

that the giant graviton as a classical solution can exist due to the presence of the constant R-R flux [18]. Then, in particular, we showed the quantum stability of the giant graviton (supersymmetric fuzzy sphere solution) and instability of the non-supersymmetric fuzzy sphere solution in Ref. [19]. As well as by our work, the classical solutions in the matrix model have been widely investigated [20]. Moreover, in Ref. [21] we also derived the less supersymmetric background with 24 supercharges from the KG solution and studied the type IIA string from the supergravity analysis and the double dimensional reduction [22]. Furthermore, we also investigated matrix string theories. The supersymmetries of the above type IIA string were initially discussed in [23].

In this paper we will continue to study the supermembrane on the pp-wave and consider the ground-state wave function of the supermembrane on the pp-wave by the use of the quantum-mechanical procedure of de Wit-Hoppe-Nicolai [11]. It is well-known that a single supermembrane in the flat space is unstable [24]. This instability is inherently based on the continuous spectrum of this system. However, the action of the supermembrane (or matrix model) on the pp-wave contains the bosonic and the fermionic mass terms and the Myers term, and hence it would be possible to expect that the spectrum of the system might be discrete or that a single supermembrane might be stabilized. Thus, it would be very interesting to study the ground state or spectrum of this system. To accomplish this, the complex-spinor notation ($Spin(7) \times U(1)$ formalism) will be introduced. We calculate the superalgebra of complex supercharges. Then, we will investigate the zero-mode Hamiltonian of the system, whose spectrum is non-trivial in contrast to the flat case. This system is identical with the abelian matrix model describing the motion of superparticles on the pp-wave. The Hamiltonian resembles the supersymmetric harmonic oscillator but the supercharges do not commute with the Hamiltonian. The supergravity multiplet **128** (boson) + **128** (fermion) in the flat space comes to split non-trivially. As a result, a unique supersymmetric ground state is left, and all the other states are lifted up. Also, we show that the zero-mode Hamiltonian is positive definite and explicitly construct the ground-state wave function for the zero-mode Hamiltonian. It is represented by the product of the ground states of the bosonic harmonic oscillators and the fermionic variables. For this ground state, both the orbital and the spin angular momenta of $SO(3) \times SO(6)$ vanish.

Furthermore, we investigate the nonzero-mode Hamiltonian and its ground-state wave function. For example, by truncating the variables, the supersymmetric ground-state wave function is explicitly constructed in the $SU(2)$ matrix model case. This ground state seems non-

normalizable, but its L^2 -norm can be represented by an asymptotic series with respect to $1/\mu^3$. The zeroth order term corresponds to the kinematical contribution to the ground state, while the corrections with the powers of $1/\mu^3$ are related to the dynamical contributions arising from non-diagonal elements of matrix variables. In the IR region, the matrix variables are restricted to the abelian part. From these considerations, it might be understood that the dynamical effect tends to spoil down the normalizability of the zero-mode ground state as the system flows from the IR region to the UV one, at least in our example.

This paper is organized as follows. In section 2 we introduce the complex-spinor notation by following the quantum-mechanical procedure of de Wit-Hoppe-Nicolai. Section 3 is devoted to the study of the zero-mode Hamiltonian of the system. We can obtain the spectrum, and construct the unique supersymmetric ground state. Also, the positive definiteness of the zero-mode Hamiltonian is shown. In section 4 we discuss the nonzero-mode Hamiltonian and construct an example of the ground-state wave function with the truncation of the variables in $N = 2$ case. Its L^2 -norm can be represented by an asymptotic series. Section 5 is devoted to conclusions and discussions. In appendix, we discuss that the ground state of the zero-mode Hamiltonian is an $SO(3) \times SO(6)$ singlet.

2 Supercharges and Superalgebra

In this section, we shall start with our setup and introduce the complex-spinor notation ($Spin(7) \times U(1)$ formalism), which is convenient to study the ground state in later sections. We will also obtain the complex representation of the superalgebra.

2.1 Setup

The Lagrangian of the supermembrane on the KG background in the light-cone gauge is written in terms of an $SO(9)$ spinor ψ as

$$\begin{aligned}
w^{-1}\mathcal{L} = & \frac{1}{2}D_\tau X^r D_\tau X^r - \frac{1}{4}(\{X^r, X^s\})^2 - \frac{1}{2}\left(\frac{\mu}{3}\right)^2 \sum_{I=1}^3 X_I^2 - \frac{1}{2}\left(\frac{\mu}{6}\right)^2 \sum_{I'=4}^9 X_{I'}^2 \\
& - \frac{\mu}{6} \sum_{I,J,K=1}^3 \epsilon_{IJK} X^K \{X^I, X^J\} + i\psi^T \gamma^r \{X^r, \psi\} + i\psi^T D_\tau \psi + i\frac{\mu}{4}\psi^T \gamma_{123} \psi. \quad (2.1)
\end{aligned}$$

Here “ τ ” is the time coordinate on the worldvolume and $\{, \}$ is Lie bracket given using an arbitrary function $w(\sigma)$ of worldvolume spatial coordinates σ^a ($a = 1, 2$)

$$\{A, B\} \equiv \frac{1}{w} \epsilon^{ab} \partial_a A \partial_b B, \quad (a, b = 1, 2).$$

with $\partial_a \equiv \frac{\partial}{\partial \sigma^a}$. Also this theory has large residual gauge symmetry called the area-preserving diffeomorphism (APD) and the covariant derivative for this gauge symmetry is defined by a gauge connection ω

$$D_\tau X^r \equiv \partial_\tau X^r - \{\omega, X^r\}. \quad (2.2)$$

The original symmetries cannot be manifestly seen in the light-cone gauge, but the Lagrangian (2.1) still has residual supersymmetries,

$$\begin{aligned} \delta_\epsilon X^r &= 2\psi^T \gamma^r \epsilon(\tau), \quad \delta_\epsilon \omega = 2\psi^T \epsilon(\tau), \\ \delta_\epsilon \psi &= -i D_\tau X^r \gamma_r \epsilon(\tau) + \frac{i}{2} \{X^r, X^s\} \gamma_{rs} \epsilon(\tau) \\ &\quad + \frac{\mu}{3} i \sum_{I=1}^3 X^I \gamma_I \gamma_{123} \epsilon(\tau) - \frac{\mu}{6} i \sum_{I'=4}^9 X^{I'} \gamma_{I'} \gamma_{123} \epsilon(\tau), \\ \epsilon(\tau) &= \exp\left(\frac{\mu}{12} \gamma_{123} \tau\right) \epsilon_0 \quad (\epsilon_0 : \text{constant spinor}). \end{aligned} \quad (2.3)$$

These transformation rules are 16 linearly-realized supersymmetries on the maximally supersymmetric pp-wave. In taking the limit $\mu \rightarrow 0$, we recover the supersymmetry transformations in the flat space. In the context of the eleven-dimensional supersymmetries, these correspond to the dynamical supersymmetries. The Lagrangian (2.1) has another 16 nonlinearly realized supersymmetries,

$$\begin{aligned} \delta_\eta X^r &= 0, \quad \delta_\eta \omega = 0, \\ \delta_\eta \psi &= \eta(\tau), \\ \eta(\tau) &= \exp\left(-\frac{\mu}{4} \gamma_{123} \tau\right) \eta_0, \quad (\eta_0 : \text{constant spinor}), \end{aligned} \quad (2.4)$$

which correspond to the kinematical supersymmetries in the eleven-dimensional theory.

In our previous work [14], we derived the supercharges and superalgebra of the supermembrane theory on the pp-wave background using the $SO(9)$ formalism. The supercharges Q^+ for the dynamical supersymmetries and Q^- for the kinematical supersymmetries are obtained as

Noether charges

$$Q^+ = \int d^2\sigma w \left[-2e^{-\frac{\mu}{12}\gamma_{123}\tau} \left(DX^r \gamma_r \psi + \frac{1}{2} \{X^r, X^s\} \gamma_{rs} \psi \right. \right. \\ \left. \left. + \frac{\mu}{3} \sum_{I=1}^3 X^I \gamma_I \gamma_{123} \psi + \frac{\mu}{6} \sum_{I'=4}^9 X^{I'} \gamma_{I'} \gamma_{123} \psi \right) \right], \quad (2.5)$$

$$Q^- = \int d^2\sigma w \left[-2ie^{\frac{\mu}{4}\gamma_{123}\tau} \psi \right] \\ = -2ie^{\frac{\mu}{4}\gamma_{123}\tau} \psi_0, \quad (2.6)$$

where ψ_0 is the zero-mode of ψ and we used the normalization with $\int d^2\sigma w(\sigma) = 1$.

By the use of the Dirac brackets,

$$\{X^r, D_\tau X_s\}_{\text{DB}} = \frac{1}{w} \delta_s^r \delta^{(2)}(\sigma - \sigma'), \quad (2.7)$$

$$\{\psi_\alpha(\sigma), \psi_\beta^T(\sigma')\}_{\text{DB}} = -\frac{i}{2w} \delta_{\alpha\beta} \delta^{(2)}(\sigma - \sigma'), \quad (2.8)$$

the superalgebra can be calculated and we obtain the following results

$$i \left\{ \frac{1}{\sqrt{2}} Q_\alpha^-, \frac{1}{\sqrt{2}} (Q^-)_\beta^T \right\}_{\text{DB}} = -\delta_{\alpha\beta}, \quad (2.9)$$

$$i \left\{ \frac{1}{\sqrt{2}} Q_\alpha^+, \frac{1}{\sqrt{2}} (Q^-)_\beta^T \right\}_{\text{DB}} = i \sum_{I=1}^3 \left[\left(P_0^I + \frac{\mu}{3} X_0^I \gamma_{123} \right) \gamma_I e^{-\frac{\mu}{3}\gamma_{123}\tau} \right]_{\alpha\beta} \quad (2.10)$$

$$+ i \sum_{I'=4}^9 \left[\left(P_0^{I'} - \frac{\mu}{6} X_0^{I'} \gamma_{123} \right) \gamma_{I'} e^{-\frac{\mu}{6}\gamma_{123}\tau} \right]_{\alpha\beta},$$

$$i \left\{ \frac{1}{\sqrt{2}} Q_\alpha^+, \frac{1}{\sqrt{2}} (Q^+)_\beta^T \right\}_{\text{DB}} = 2H \delta_{\alpha\beta} \quad (2.11)$$

$$+ \frac{\mu}{3} \sum_{I,J=1}^3 M_0^{IJ} (\gamma_{IJ} \gamma_{123})_{\alpha\beta} - \frac{\mu}{6} \sum_{I',J'=4}^9 M_0^{I'J'} (\gamma_{I'J'} \gamma_{123})_{\alpha\beta} \\ - 2 \sum_{I=1}^3 \int d^2\sigma \varphi X_I (\gamma^I)_{\alpha\beta} - 2 \sum_{I'=4}^9 \int d^2\sigma \varphi X_{I'} \left(\gamma^{I'} e^{\frac{\mu}{6}\gamma_{123}\tau} \right)_{\alpha\beta},$$

where we omitted the central charges since these are not discussed in this paper. Here φ describes the constraint condition,

$$\varphi = w \{w^{-1} P^r, X^r\} + iw \{\psi^T, \psi\}. \quad (2.12)$$

The $P^r \equiv w D_\tau X^r$ and $S_\alpha = iw \psi_\alpha^T$ are canonical momenta of the X^r and ψ , respectively. The M^{IJ} and $M^{I'J'}$ are defined by

$$M^{IJ} = X^I P^J - P^I X^J - \frac{1}{2} S \gamma^{IJ} \psi, \quad (2.13)$$

$$M^{I'J'} = X^{I'} P^{J'} - P^{I'} X^{J'} - \frac{1}{2} S \gamma^{I'J'} \psi, \quad (2.14)$$

and the $SO(3) \times SO(6)$ Lorentz generators M_0^{IJ} and $M_0^{I'J'}$ are given as

$$M_0^{IJ} \equiv \int d^2 \sigma M^{IJ}, \quad (2.15)$$

$$M_0^{I'J'} \equiv \int d^2 \sigma M^{I'J'}. \quad (2.16)$$

They satisfy the $SO(3) \times SO(6)$ Lorentz algebra,

$$\{M_0^{IJ}, M_0^{KL}\}_{\text{DB}} = \delta^{IK} M_0^{JL} - \delta^{IL} M_0^{JK} - \delta^{JK} M_0^{IL} + \delta^{JL} M_0^{IK}, \quad (2.17)$$

$$\{M_0^{I'J'}, M_0^{K'L'}\}_{\text{DB}} = \delta^{I'K'} M_0^{J'L'} - \delta^{I'L'} M_0^{J'K'} - \delta^{J'K'} M_0^{I'L'} + \delta^{J'L'} M_0^{I'K'}. \quad (2.18)$$

The zero-modes of $P^r (\equiv w D_\tau X^r)$ and X^r are written by

$$P_0^r \equiv \int d^2 \sigma w D_\tau X^r, \quad X_0^r \equiv \int d^2 \sigma w X^r. \quad (2.19)$$

Also, the Hamiltonian H is given by

$$\begin{aligned} H = \int d^2 \sigma w & \left[\frac{1}{2} \left(\frac{P^r}{w} \right)^2 + \frac{1}{4} \{X^r, X^s\}^2 + \frac{1}{2} \left(\frac{\mu}{3} \right)^2 \sum_{I=1}^3 (X^I)^2 + \frac{1}{2} \left(\frac{\mu}{6} \right)^2 \sum_{I'=4}^9 (X^{I'})^2 \right. \\ & \left. + \frac{\mu}{6} \sum_{I,J,K=1}^3 \epsilon_{IJK} X^K \{X^I, X^J\} - w^{-1} \frac{\mu}{4} S \gamma_{123} \psi - w^{-1} S \gamma_r \{X^r, \psi\} \right]. \end{aligned} \quad (2.20)$$

In the next section, we will introduce the complex spinor notation in order to study the ground-state wave function.

2.2 $Spin(7) \times U(1)$ Formalism

Here we shall introduce the $Spin(7) \times U(1)$ formalism following Ref. [11]. Hereafter, we use the decomposition of the $SO(9)$ gamma matrices into the $SO(7)$ ones as follows:

$$\begin{aligned} \gamma_i &= \Gamma_i \otimes \sigma_2 = \begin{pmatrix} 0 & -i\Gamma_i \\ i\Gamma_i & 0 \end{pmatrix}, \quad (i = 1, \dots, 7), \\ \gamma_8 &= \mathbf{1}_8 \otimes \sigma_1 = \begin{pmatrix} 0 & \mathbf{1}_8 \\ \mathbf{1}_8 & 0 \end{pmatrix}, \quad \gamma_9 = \mathbf{1}_8 \otimes \sigma_3 = \begin{pmatrix} \mathbf{1}_8 & 0 \\ 0 & -\mathbf{1}_8 \end{pmatrix}. \end{aligned} \quad (2.21)$$

The above $SO(7)$ gamma matrices satisfy the following relations:

$$(\Gamma_i)^\dagger = \Gamma_i, \quad (\Gamma_i)^T = -\Gamma_i, \quad (\Gamma_i)^* = -\Gamma_i. \quad (2.22)$$

To begin, let us decompose the real 16-component $SO(9)$ spinor into two real 8-component spinors $\psi^{(+)}$ and $\psi^{(-)}$, which are eigenspinors for the chirality matrix γ_9 defined by

$$\gamma_9 \psi^{(\pm)} = (\pm 1) \cdot \psi^{(\pm)}. \quad (2.23)$$

In this time we can express eigenspinors as $\psi_\alpha^{(+)} = \psi_\alpha$ and $\psi_\beta^{(-)} = \psi_{8+\beta}$ ($\alpha, \beta = 1, \dots, 8$). Here, we shall introduce a complex spinor λ defined by

$$\lambda_\alpha = \psi_\alpha^{(+)} + i\psi_\alpha^{(-)}, \quad (\alpha = 1, \dots, 8), \quad (2.24)$$

that is, we express the real 16-component $SO(9)$ spinor as a single 8-component complex spinor λ . This complex spinor is linearly-realized under the $SO(7) \times U(1)$ transformation, which is the subgroup of the original $SO(9)$. The following expression

$$\psi = \begin{pmatrix} \psi^{(+)} \\ \psi^{(-)} \end{pmatrix} = \frac{1}{2} \begin{pmatrix} \lambda + \lambda^\dagger \\ -i(\lambda - \lambda^\dagger) \end{pmatrix} \quad (2.25)$$

is also useful for rewriting the supercharges.

The bosonic coordinates X^8 and X^9 are combined into a complex coordinate

$$Z = \frac{1}{\sqrt{2}}(X^8 + iX^9), \quad (2.26)$$

which transforms linearly under the $U(1)$ subgroup. The canonical momentum for Z is given by

$$\mathcal{P} = \frac{1}{\sqrt{2}}(P^8 - iP^9). \quad (2.27)$$

By the standard procedure, the Dirac brackets for these variables are given by

$$\begin{aligned} \{X^i(\tau, \sigma), P^j(\tau, \sigma')\}_{\text{DB}} &= \delta^{ij} \delta^{(2)}(\sigma - \sigma'), \quad (i, j = 1, \dots, 7), \\ \{Z(\tau, \sigma), \mathcal{P}(\tau, \sigma')\}_{\text{DB}} &= \delta^{(2)}(\sigma - \sigma'), \\ \{\lambda_\alpha(\tau, \sigma), \lambda_\beta^\dagger(\tau, \sigma')\}_{\text{DB}} &= -\frac{i}{w} \delta_{\alpha\beta} \delta^{(2)}(\sigma - \sigma'), \quad (\alpha, \beta = 1, \dots, 8). \end{aligned} \quad (2.28)$$

Next, let us rewrite the supercharge Q^+ in terms of the complex spinor λ . We decompose the supercharge Q^+ into $(Q^+)^{(\pm)}$ as

$$Q^+ = \begin{pmatrix} (Q^+)^{(+)} \\ (Q^+)^{(-)} \end{pmatrix},$$

and define the complex supercharge Q by

$$Q \equiv \frac{1}{2} ((Q^+)^{(+)} + i(Q^+)^{(-)}) . \quad (2.29)$$

Thus, after straightforward calculation, we can obtain the expression of the complex supercharge as

$$\begin{aligned} Q = & \int d^2\sigma e^{\frac{\mu}{12}\Gamma_{123}\tau} \left[\left(P^i \Gamma_i - \frac{1}{2} w \{X^i, X^j\} \Gamma_{ij} + w \{Z, \bar{Z}\} \right. \right. \\ & - \frac{\mu}{3} w \sum_{I=1}^3 X^I \Gamma_I \Gamma_{123} - \frac{\mu}{6} w \sum_{m=4}^7 X^m \Gamma_m \Gamma_{123} \Big) \lambda \\ & \left. \left. - \sqrt{2} i \left(\mathcal{P} - w \{X^i, \bar{Z}\} \Gamma_i + \frac{\mu}{6} w \bar{Z} \Gamma_{123} \right) \lambda^\dagger \right] , \end{aligned} \quad (2.30)$$

and its complex conjugate is given by

$$\begin{aligned} Q^\dagger = & - \int d^2\sigma e^{-\frac{\mu}{12}\Gamma_{123}\tau} \left[\left(P^i \Gamma_i + \frac{1}{2} w \{X^i, X^j\} \Gamma_{ij} + w \{Z, \bar{Z}\} \right. \right. \\ & + \frac{\mu}{3} w \sum_{I=1}^3 X^I \Gamma_I \Gamma_{123} + \frac{\mu}{6} w \sum_{m=4}^7 X^m \Gamma_m \Gamma_{123} \Big) \lambda^\dagger \\ & \left. \left. - \sqrt{2} i \left(\bar{\mathcal{P}} + w \{X^i, Z\} \Gamma_i - \frac{\mu}{6} w Z \Gamma_{123} \right) \lambda \right] . \end{aligned} \quad (2.31)$$

The superalgebra of these complex supercharges is computed as follows:

$$i\{Q_\alpha, Q_\beta\}_{\text{DB}} = i\frac{\mu}{3} \sum_{m=4}^7 M_0^{m+} (\Gamma_m \Gamma_{123})_{\alpha\beta} - 2\sqrt{2}i \int d^2\sigma \varphi \bar{Z} \left(e^{\frac{\mu}{6}\Gamma_{123}\tau} \right)_{\alpha\beta} , \quad (2.32)$$

$$i\{Q_\alpha^\dagger, Q_\beta^\dagger\}_{\text{DB}} = -i\frac{\mu}{3} \sum_{m=4}^7 M_0^{m-} (\Gamma_m \Gamma_{123})_{\alpha\beta} + 2\sqrt{2}i \int d^2\sigma \varphi Z \left(e^{-\frac{\mu}{6}\Gamma_{123}\tau} \right)_{\alpha\beta} , \quad (2.33)$$

$$\begin{aligned} i\{Q_\alpha, Q_\beta^\dagger\}_{\text{DB}} = & 2H\delta_{\alpha\beta} + i\frac{\mu}{3} M_0^{89} (\Gamma_{123})_{\alpha\beta} - \frac{\mu}{3} \sum_{I,J=1}^3 M_0^{IJ} (\Gamma_{IJ} \Gamma_{123})_{\alpha\beta} \\ & + \frac{\mu}{6} \sum_{m,n=4}^7 M_0^{mn} (\Gamma_{mn} \Gamma_{123})_{\alpha\beta} + 2 \sum_{I=1}^3 \int d^2\sigma \varphi X_I (\Gamma_I)_{\alpha\beta} \\ & + 2 \sum_{m=4}^7 \int d^2\sigma \varphi X_m \left(\Gamma_m e^{-\frac{\mu}{6}\Gamma_{123}\tau} \right)_{\alpha\beta} , \end{aligned} \quad (2.34)$$

where the zero-modes of angular momenta M_0^{IJ} , M_0^{mn} , $M_0^{m\pm}$ and M_0^{89} are rewritten in terms

of the complex notation as follows:

$$M_0^{IJ} = \int d^2\sigma M^{IJ}, \quad M^{IJ} = X^I P^J - X^J P^I - iw \frac{1}{2} \lambda^\dagger \Gamma^{IJ} \lambda, \quad (2.35)$$

$$M_0^{mn} = \int d^2\sigma M^{mn}, \quad M^{mn} = X^m P^n - X^n P^m - iw \frac{1}{2} \lambda^\dagger \Gamma^{mn} \lambda, \quad (2.36)$$

$$M_0^{m+} = \int d^2\sigma M^{m+}, \quad M^{m+} = \sqrt{2} (X^m \mathcal{P} - P^m \bar{Z}) - \frac{1}{2} w \lambda \Gamma_m \lambda, \quad (2.37)$$

$$M_0^{m-} = \int d^2\sigma M^{m-}, \quad M^{m-} = \sqrt{2} (X^m \bar{\mathcal{P}} - P^m Z) - \frac{1}{2} w \lambda^\dagger \Gamma_m \lambda^\dagger, \quad (2.38)$$

$$M_0^{89} = \int d^2\sigma M^{89}, \quad M^{89} = i (Z \mathcal{P} - \bar{Z} \bar{\mathcal{P}}) + \frac{1}{4} w (\lambda^\dagger \lambda - \lambda \lambda^\dagger). \quad (2.39)$$

In the above superalgebra, the Hamiltonian H is given by

$$\begin{aligned} H = & \int d^2\sigma w \left[\frac{1}{2} \left(\frac{P^i}{w} \right)^2 + w^{-2} |\mathcal{P}|^2 + \frac{1}{4} \{X^i, X^j\}^2 + |\{X^i, Z\}|^2 + \frac{1}{2} |\{Z, \bar{Z}\}|^2 \right. \\ & + \frac{1}{2} \left(\frac{\mu}{3} \right)^2 \sum_{I=1}^3 (X^I)^2 + \frac{1}{2} \left(\frac{\mu}{6} \right)^2 \sum_{m=4}^7 (X^m)^2 + \left(\frac{\mu}{6} \right)^2 Z \bar{Z} + \frac{\mu}{6} \sum_{I,J,K=1}^3 \epsilon_{IJK} X^K \{X^I, X^J\} \\ & \left. - i \lambda \Gamma_i \{X^i, \lambda^\dagger\} - \frac{1}{\sqrt{2}} (\lambda \{Z, \lambda\} - \lambda^\dagger \{\bar{Z}, \lambda^\dagger\}) - i \frac{\mu}{4} \lambda \Gamma_{123} \lambda^\dagger \right]. \end{aligned} \quad (2.40)$$

The constraint condition is also expressed by

$$w^{-1} \varphi = \{w^{-1} P^i, X^i\} + \{w^{-1} \mathcal{P}, Z\} + \{w^{-1} \bar{\mathcal{P}}, \bar{Z}\} + i \{\lambda, \lambda^\dagger\}. \quad (2.41)$$

When we contract the spinor indices α and β , the Hamiltonian is represented with the above supercharges as

$$H = \frac{1}{16} i \{Q_\alpha, Q_\alpha^\dagger\}_{\text{DB}}. \quad (2.42)$$

Now, we shall move to a matrix representation by replacing the variables with matrices, according to:

$$\begin{aligned} X(\xi^i) & \longrightarrow X(\tau), \\ \psi(\xi^i) & \longrightarrow \psi(\tau), \\ \int d^2\sigma w(\sigma) & \longrightarrow N \text{Tr}, \\ \{, \} & \longrightarrow -i[,]. \end{aligned}$$

In this case, the Hamiltonian is rewritten into

$$\begin{aligned}
H = N \text{Tr} & \left[\frac{1}{2} (P^i)^2 + \mathcal{P} \bar{\mathcal{P}} - \frac{1}{4} [X^i, X^j]^2 + |[X^i, Z]|^2 + \frac{1}{2} |[Z, \bar{Z}]|^2 \right. \\
& + \frac{1}{2} \left(\frac{\mu}{3} \right)^2 \sum_{I=1}^3 (X^I)^2 + \frac{1}{2} \left(\frac{\mu}{6} \right)^2 \sum_{m=4}^7 (X^m)^2 + \left(\frac{\mu}{6} \right)^2 Z \bar{Z} - i \frac{\mu}{3} \sum_{I,J,K=1}^3 \epsilon_{IJK} X^I X^J X^K \\
& \left. - \lambda \Gamma_i [X^i, \lambda^\dagger] + \frac{1}{\sqrt{2}} i (\lambda [Z, \lambda] - \lambda^\dagger [\bar{Z}, \lambda^\dagger]) - i \frac{\mu}{4} \lambda \Gamma_{123} \lambda^\dagger \right], \tag{2.43}
\end{aligned}$$

where the canonical momenta are redefined by $P^i \equiv D_\tau X^i$, $\mathcal{P} \equiv D_\tau \bar{Z}$, $\bar{\mathcal{P}} \equiv D_\tau Z$. The variables are expressed by the $U(N)$ generators T^A ($A = 0, 1, \dots, N^2 - 1$, $T^0 = \mathbf{1}$) as follows:*

$$X^i = X_0^i \mathbf{1} + \sum_{A=1}^{N^2-1} X_A^i T^A, \quad P^i = P_0^i \mathbf{1} + \sum_{A=1}^{N^2-1} P_A^i T^A, \tag{2.44}$$

$$Z = Z_0 \mathbf{1} + \sum_{A=1}^{N^2-1} Z_A T^A, \quad \mathcal{P} = \mathcal{P}_0 \mathbf{1} + \sum_{A=1}^{N^2-1} \mathcal{P}_A T^A, \tag{2.45}$$

$$\lambda_\alpha = \lambda_{0\alpha} \mathbf{1} + \sum_{A=1}^{N^2-1} \lambda_{A\alpha} T^A, \quad \lambda_\alpha^\dagger = \lambda_{0\alpha}^\dagger \mathbf{1} + \sum_{A=1}^{N^2-1} \lambda_{A\alpha}^\dagger T^A. \tag{2.46}$$

Also, the $U(N)$ generators T^A ($A = 0, 1, \dots, N^2 - 1$) satisfy the following relations:

$$[T^A, T^B] = i f_{ABC} T^C, \quad \text{Tr}(T^A T^B) = \frac{1}{N} \delta^{AB}.$$

where the structure constant f_{ABC} is a completely antisymmetric tensor satisfying $f_{0AB} = 0$.

Now, the Dirac brackets are replaced with the following (anti-)commutation relations

$$\begin{aligned}
[X_A^i, P_B^j] &= i \delta^{ij} \delta_{AB}, \quad (i, j = 1, \dots, 7), \\
[Z_A, \mathcal{P}_B] &= [\bar{Z}_A, \bar{\mathcal{P}}_B] = i \delta_{AB}, \\
\{\lambda_{A\alpha}, \lambda_{B\beta}^\dagger\} &= \delta_{\alpha\beta} \delta_{AB}, \quad (\alpha, \beta = 1, \dots, 8).
\end{aligned} \tag{2.47}$$

Therefore, we can represent the conjugate momenta by differential operators

$$\begin{aligned}
P_A^i &= -i \frac{\partial}{\partial X_A^i}, \quad \mathcal{P}^A = -i \frac{\partial}{\partial Z_A}, \\
\bar{\mathcal{P}}^A &= -i \frac{\partial}{\partial \bar{Z}_A}, \quad \lambda_{A\alpha}^\dagger = \frac{\partial}{\partial \lambda_{A\alpha}}.
\end{aligned} \tag{2.48}$$

*We assume that the topology of the membrane is sphere.

Then we obtain the following expressions for supercharges as

$$\begin{aligned}
Q_\alpha = & \left(e^{\frac{\mu}{12}\Gamma_{123}\tau} \right)_{\alpha\beta} \left[\left(-i(\Gamma_i)_{\beta\gamma} \frac{\partial}{\partial X_A^i} - \frac{1}{2} f_{ABC} X_B^i X_C^j (\Gamma_{ij})_{\beta\gamma} + f_{ABC} Z^B \bar{Z}^C \delta_{\beta\gamma} \right. \right. \\
& - \frac{\mu}{3} \sum_{I=1}^3 X_A^I (\Gamma_I \Gamma_{123})_{\beta\gamma} - \frac{\mu}{6} \sum_{m=4}^7 X_A^m (\Gamma_m \Gamma_{123})_{\beta\gamma} \Big) \lambda_{A\gamma} \\
& \left. - \sqrt{2} \left(\frac{\partial}{\partial Z_A} \delta_{\beta\gamma} - i f_{ABC} X_B^i \bar{Z}_C (\Gamma_i)_{\beta\gamma} + i \frac{\mu}{6} \bar{Z}_A (\Gamma_{123})_{\beta\gamma} \right) \frac{\partial}{\partial \lambda_{A\gamma}} \right], \quad (2.49)
\end{aligned}$$

and its complex conjugate is given by

$$\begin{aligned}
Q_\alpha^\dagger = & - \left(e^{-\frac{\mu}{12}\Gamma_{123}\tau} \right)_{\alpha\beta} \left[\left(-i(\Gamma_i)_{\beta\gamma} \frac{\partial}{\partial X_A^i} + \frac{1}{2} f_{ABC} X_B^i X_C^j (\Gamma_{ij})_{\beta\gamma} + f_{ABC} Z_B \bar{Z}_C \delta_{\beta\gamma} \right. \right. \\
& + \frac{\mu}{3} \sum_{I=1}^3 X_A^I (\Gamma_I \Gamma_{123})_{\beta\gamma} + \frac{\mu}{6} \sum_{m=4}^7 X_A^m (\Gamma_m \Gamma_{123})_{\beta\gamma} \Big) \frac{\partial}{\partial \lambda_{A\gamma}} \\
& \left. - \sqrt{2} \left(\frac{\partial}{\partial \bar{Z}_A} \delta_{\beta\gamma} + i f_{ABC} X_B^i Z_C (\Gamma_i)_{\beta\gamma} - i \frac{\mu}{6} Z_A (\Gamma_{123})_{\beta\gamma} \right) \lambda_{A\gamma} \right]. \quad (2.50)
\end{aligned}$$

In this representation, the superalgebra becomes

$$\{Q_\alpha, Q_\beta\} = i \frac{\mu}{3} \sum_{m=4}^7 M_0^{m+} (\Gamma_m \Gamma_{123})_{\alpha\beta} - 2\sqrt{2} i \varphi_A \bar{Z}_A \left(e^{\frac{\mu}{6}\Gamma_{123}\tau} \right)_{\alpha\beta}, \quad (2.51)$$

$$\{Q_\alpha^\dagger, Q_\beta^\dagger\} = -i \frac{\mu}{3} \sum_{m=4}^7 M_0^{m-} (\Gamma_m \Gamma_{123})_{\alpha\beta} + 2\sqrt{2} i \varphi_A Z_A \left(e^{-\frac{\mu}{6}\Gamma_{123}\tau} \right)_{\alpha\beta}, \quad (2.52)$$

$$\begin{aligned}
\{Q_\alpha, Q_\beta^\dagger\} = & 2H \delta_{\alpha\beta} - \frac{\mu}{3} \sum_{I,J=1}^3 M_0^{IJ} (\Gamma_{IJ} \Gamma_{123})_{\alpha\beta} + \frac{\mu}{6} \sum_{m,n=4}^7 M_0^{mn} (\Gamma_{mn} \Gamma_{123})_{\alpha\beta} \\
& + i \frac{\mu}{3} M_0^{89} (\Gamma_{123})_{\alpha\beta} + 2 \sum_{I=1}^3 \varphi_A X_A^I (\Gamma_I)_{\alpha\beta} + 2 \sum_{m=4}^7 \varphi_A X_A^m \left(\Gamma_m e^{-\frac{\mu}{6}\Gamma_{123}\tau} \right)_{\alpha\beta}, \quad (2.53)
\end{aligned}$$

where the zero-modes of angular momenta are represented by

$$M_0^{IJ} = -i X_A^I \frac{\partial}{\partial X_A^J} + i X_A^J \frac{\partial}{\partial X_A^I} - \frac{i}{2} \lambda_A \Gamma^{IJ} \frac{\partial}{\partial \lambda_A}, \quad (2.54)$$

$$M_0^{mn} = -i X_A^m \frac{\partial}{\partial X_A^n} + i X_A^n \frac{\partial}{\partial X_A^m} - \frac{i}{2} \lambda_A \Gamma^{mn} \frac{\partial}{\partial \lambda_A}, \quad (2.55)$$

$$M_0^{m+} = -\sqrt{2} i \left(X_A^m \frac{\partial}{\partial Z_A} - \bar{Z}_A \frac{\partial}{\partial X_A^m} \right) - \frac{1}{2} \lambda_A \Gamma^m \lambda_A, \quad (2.56)$$

$$M_0^{m-} = -\sqrt{2} i \left(X_A^m \frac{\partial}{\partial \bar{Z}_A} - \frac{\partial}{\partial X_A^m} Z_A \right) - \frac{1}{2} \frac{\partial}{\partial \lambda_A} \Gamma^m \frac{\partial}{\partial \lambda_A}, \quad (2.57)$$

$$M_0^{89} = Z_A \frac{\partial}{\partial Z_A} - \bar{Z}_A \frac{\partial}{\partial \bar{Z}_A} - \frac{1}{2} \lambda_A \frac{\partial}{\partial \lambda_A} + 2 \cdot \dim U(N). \quad (2.58)$$

and the constraint condition φ_A is expressed by

$$\varphi_A \equiv if_{ABC} \left(X_B^i \frac{\partial}{\partial X_C^i} + Z_B \frac{\partial}{\partial Z_C} + \bar{Z}_B \frac{\partial}{\partial \bar{Z}_C} + \lambda_B \frac{\partial}{\partial \lambda_C} \right). \quad (2.59)$$

It is assumed that physical states should satisfy the constraint condition i.e., $\varphi\Psi_{\text{phys}} = 0$. Our purpose is to study the ground state. Thus, we shall also restrict ourselves to the $SO(3) \times SO(6)$ singlet satisfying

$$M^{IJ}\Psi_{\text{phys}} = M^{mn}\Psi_{\text{phys}} = M^{m\pm}\Psi_{\text{phys}} = M^{89}\Psi_{\text{phys}} = 0, \quad (2.60)$$

since the ground state should have no angular momenta. The discussion on the states with nonzero angular momenta would be also possible, but we do not consider them in this paper.

Furthermore, the associated Hamiltonian is written down as

$$\begin{aligned} H &= \frac{1}{16} \{Q_\alpha, Q_\alpha^\dagger\} \\ &= -\frac{1}{2} \frac{\partial^2}{\partial X_A^i \partial X_A^i} - \frac{\partial^2}{\partial Z_A \partial \bar{Z}_A} + \frac{1}{4} f_{ABC} f_{ADE} X_B^i X_C^j X_D^i X_E^j \\ &\quad + f_{ABC} f_{ADE} X_B^i Z_C X_D^i \bar{Z}_E + \frac{1}{2} f_{ABC} f_{ADE} Z_B \bar{Z}_C \bar{Z}_D Z_E \\ &\quad + \frac{1}{2} \left(\frac{\mu}{3}\right)^2 \sum_{I=1}^3 (X_A^I)^2 + \frac{1}{2} \left(\frac{\mu}{6}\right)^2 \sum_{m=4}^7 (X_A^m)^2 + \left(\frac{\mu}{6}\right)^2 Z_A \bar{Z}_A + \frac{\mu}{6} \sum_{I,J,K=1}^3 \epsilon_{IJK} f_{ABC} X_A^I X_B^J X_C^K \\ &\quad + if_{ABC} X_A^i \lambda_B \Gamma_i \frac{\partial}{\partial \lambda_C} + \frac{1}{\sqrt{2}} f_{ABC} \left(Z_A \lambda_B \lambda_C - \bar{Z}_A \frac{\partial}{\partial \lambda_B} \frac{\partial}{\partial \lambda_C} \right) - i \frac{\mu}{4} \lambda_A \Gamma_{123} \frac{\partial}{\partial \lambda_A}. \end{aligned} \quad (2.61)$$

Hereafter, by the use of the supercharges, the Hamiltonian, the angular momenta and the constraint condition, we will study the ground-state wave function analytically.

3 Supersymmetric Quantum Mechanics of Zero-mode Hamiltonian

In this section, we shall consider the zero-mode ground-state wave function. We can extract the zero-mode part of the dynamical supercharges and the results are given by

$$\begin{aligned} Q_\alpha^{(0)} &= \left(e^{\frac{\mu}{12} \Gamma_{123} \tau} \right)_{\alpha\beta} \left[\left(-i(\Gamma_i)_{\beta\gamma} \frac{\partial}{\partial X_0^i} - \frac{\mu}{3} \sum_{I=1}^3 X_0^I (\Gamma_I \Gamma_{123})_{\beta\gamma} \right. \right. \\ &\quad \left. \left. - \frac{\mu}{6} \sum_{m=4}^7 X_0^m (\Gamma_m \Gamma_{123})_{\beta\gamma} \right) \lambda_{0\gamma} - \sqrt{2} \left(\frac{\partial}{\partial Z_0} \delta_{\beta\gamma} + i \frac{\mu}{6} \bar{Z}_0 (\Gamma_{123})_{\beta\gamma} \right) \frac{\partial}{\partial \lambda_{0\gamma}} \right], \quad (3.1) \end{aligned}$$

and

$$Q_\alpha^{(0)\dagger} = - \left(e^{-\frac{\mu}{12} \Gamma_{123} \tau} \right)_{\alpha\beta} \left[\left(-i(\Gamma_i)_{\beta\gamma} \frac{\partial}{\partial X_0^i} + \frac{\mu}{3} \sum_{I=1}^3 X_0^I (\Gamma_I \Gamma_{123})_{\beta\gamma} \right. \right. \\ \left. \left. + \frac{\mu}{6} \sum_{m=4}^7 X_0^m (\Gamma_m \Gamma_{123})_{\beta\gamma} \right) \frac{\partial}{\partial \lambda_{0\gamma}} - \sqrt{2} \left(\frac{\partial}{\partial \bar{Z}_0} \delta_{\beta\gamma} - i \frac{\mu}{6} Z_0 (\Gamma_{123})_{\beta\gamma} \right) \lambda_{0\gamma} \right]. \quad (3.2)$$

The zero-mode Hamiltonian $H^{(0)}$ is obtained as

$$H^{(0)} = \frac{1}{16} \{Q_\alpha^{(0)}, Q_\alpha^{(0)\dagger}\} \\ = H_B^{(0)} + H_F^{(0)}, \quad (3.3)$$

where we defined the bosonic and fermionic Hamiltonians $H_B^{(0)}$ and $H_F^{(0)}$ by

$$H_B^{(0)} \equiv -\frac{1}{2} \frac{\partial^2}{\partial X_0^i \partial X_0^i} - \frac{\partial^2}{\partial Z_0 \partial \bar{Z}_0} \\ + \frac{1}{2} \left(\frac{\mu}{3} \right)^2 \sum_{I=1}^3 (X_0^I)^2 + \frac{1}{2} \left(\frac{\mu}{6} \right)^2 \sum_{m=4}^7 (X_0^m)^2 + \left(\frac{\mu}{6} \right)^2 |Z_0|^2, \quad (3.4)$$

$$H_F^{(0)} \equiv -i \frac{\mu}{4} \lambda_{0\alpha} (\Gamma_{123})_{\alpha\beta} \frac{\partial}{\partial \lambda_{0\beta}}. \quad (3.5)$$

It is noted that the fermionic Hamiltonian for zero-modes is non-zero in the pp-wave case. Therefore, the nature of the supergravity multiplet is not simple and contains non-trivial results. The system described by the above zero-mode Hamiltonian resembles supersymmetric harmonic oscillator, although the supercharges do not commute with the Hamiltonian[†].

Here, let us study the spectrum of this system quantum mechanically. To begin, we decompose the wave function as $\Psi_B^{(0)} \otimes \Psi_F^{(0)}$. We would like to study the states of the system. We have to study the energy eigenvalue problem

$$H^{(0)} \Psi^{(0)} = (H_B^{(0)} \Psi_B^{(0)}) \otimes \Psi_F^{(0)} + \Psi_B^{(0)} \otimes (H_F^{(0)} \Psi_F^{(0)}) \\ = (E_B + E_F) \Psi^{(0)} = E \Psi^{(0)}. \quad (3.6)$$

Let us consider below the bosonic and the fermionic sectors, separately.

To begin, we would like to consider the eigenvalue problem for the bosonic zero-mode Hamiltonian (3.4). The condition $H_B^{(0)} \Psi_B^{(0)} = E_B \Psi_B^{(0)}$ can be easily solved since the bosonic Hamiltonian describes a collection of bosonic harmonic oscillators.

[†] This system has already appeared in Ref. [8] as the Lagrangian for superparticles.

Now, we shall introduce annihilation operators

$$a_0^I = \frac{i}{\sqrt{2}} \sqrt{\frac{3}{|\mu|}} \left(P_0^I - i \frac{\mu}{3} X_0^I \right), \quad (I = 1, 2, 3), \quad (3.7)$$

$$a_0^m = \frac{i}{\sqrt{2}} \sqrt{\frac{6}{|\mu|}} \left(P_0^m - i \frac{\mu}{6} X_0^m \right), \quad (m = 4, \dots, 7), \quad (3.8)$$

$$b_0 = \frac{i}{\sqrt{2}} \sqrt{\frac{6}{|\mu|}} \left(\bar{\mathcal{P}}_0 - i \frac{\mu}{6} Z_0 \right), \quad (3.9)$$

and creation operators

$$a_0^{I\dagger} = -\frac{i}{\sqrt{2}} \sqrt{\frac{3}{|\mu|}} \left(P_0^I + i \frac{\mu}{3} X_0^I \right), \quad (I = 1, 2, 3), \quad (3.10)$$

$$a_0^{m\dagger} = -\frac{i}{\sqrt{2}} \sqrt{\frac{6}{|\mu|}} \left(P_0^m + i \frac{\mu}{6} X_0^m \right), \quad (m = 4, \dots, 7), \quad (3.11)$$

$$b_0^\dagger = -\frac{i}{\sqrt{2}} \sqrt{\frac{6}{|\mu|}} \left(\mathcal{P}_0 + i \frac{\mu}{6} \bar{Z}_0 \right). \quad (3.12)$$

These operators satisfy the following relations,

$$[a_0^I, a_0^{J\dagger}] = \delta^{IJ}, \quad [a_0^m, a_0^{n\dagger}] = \delta^{mn}, \quad [b_0, b_0^\dagger] = 1. \quad (3.13)$$

By the use of the annihilation and creation operators, the bosonic zero-mode Hamiltonian $H_B^{(0)}$ can be rewritten as

$$H_B^{(0)} = \frac{|\mu|}{3} \sum_{I=1}^3 a_0^{I\dagger} a_0^I + \frac{|\mu|}{6} \left(\sum_{m=4}^7 a_0^{m\dagger} a_0^m + 2b_0^\dagger b_0 \right) + |\mu|, \quad (3.14)$$

where we note that the bosonic energy does not become zero due to the presence of the zero-point energy. This zero-point energy plays an important role to ensure the positive definiteness of the total zero-mode Hamiltonian, as we will see in the study of the fermionic sector.

As usual treatment, we can obtain all eigenstates and eigenvalues. The wave functions are given as the products of Hermite polynomials. In particular, the ground-state wave function is given by

$$\begin{aligned} \Psi_{B0}^{(0)} &= \left(\frac{|\mu|}{3\pi} \right)^{3/4} \left(\frac{|\mu|}{6\pi} \right)^{3/2} \\ &\times \exp \left[-\frac{1}{2} \cdot \frac{|\mu|}{3} \sum_{I=1}^3 (X_0^I)^2 - \frac{1}{2} \cdot \frac{|\mu|}{6} \sum_{m=4}^7 (X_0^m)^2 - \frac{|\mu|}{6} |Z_0|^2 \right], \end{aligned} \quad (3.15)$$

whose energy eigenvalue is $|\mu|$, that is,

$$H_B^{(0)} \Psi_{B0}^{(0)} = |\mu| \Psi_{B0}^{(0)}.$$

Thus, the bosonic zero-mode Hamiltonian has a lower bound as follows:

$$H_B^{(0)} \geq |\mu|. \quad (3.16)$$

Next, we shall consider the fermionic zero-mode Hamiltonian (3.5). Here, we shall decompose the complex spinor λ as

$$\begin{aligned} \lambda &= \left(\frac{1 + i\Gamma_{123}}{2} \right) \lambda + \left(\frac{1 - i\Gamma_{123}}{2} \right) \lambda \\ &\equiv \lambda^\uparrow + \lambda^\downarrow. \end{aligned} \quad (3.17)$$

By definition, $\lambda^{\uparrow,\downarrow}$ should satisfy

$$i\Gamma_{123} \lambda^{\uparrow,\downarrow} = \pm \lambda^{\uparrow,\downarrow}. \quad (3.18)$$

Now, we can rewrite the fermionic Hamiltonian as

$$H_F^{(0)} = -\frac{\mu}{4} \left(\lambda_{0\alpha}^\uparrow \frac{\partial}{\partial \lambda_{0\alpha}^\uparrow} - \lambda_{0\alpha}^\downarrow \frac{\partial}{\partial \lambda_{0\alpha}^\downarrow} \right). \quad (3.19)$$

Let us consider the zero-energy condition $H_F^{(0)} \Psi_F^{(0)} = 0$. For example, $\Psi_{F,0}^{(0)} = \text{const.}$ is a trivial solution. A pair of $\lambda^\uparrow$ and $\lambda^\downarrow$ is also a solution

$$\Psi_{F,0}^{(0)} = \lambda_{0\alpha}^\uparrow \lambda_{0\beta}^\downarrow.$$

Similarly, the following states

$$\lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow, \quad \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow \lambda_{0\alpha_3}^\uparrow \lambda_{0\beta_3}^\downarrow, \quad \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow \lambda_{0\alpha_3}^\uparrow \lambda_{0\beta_3}^\downarrow \lambda_{0\alpha_4}^\uparrow \lambda_{0\beta_4}^\downarrow, \quad (3.20)$$

also represent zero-energy states. When we note that 4 components of $\lambda_\alpha^\uparrow$ or $\lambda_\alpha^\downarrow$ are non-zero, the total number of the zero-energy states is $1 + 16 + 36 + 16 + 1 = 70$. As a result, we can represent the zero-energy ground-state wave function of the fermionic zero-mode Hamiltonian as the linear combination of these states

$$\begin{aligned} \Psi_{F,0}^{(0)} &= c_0 + c_2 \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow + c_4 \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow + c_6 \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow \lambda_{0\alpha_3}^\uparrow \lambda_{0\beta_3}^\downarrow \\ &\quad + c_8 \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow \lambda_{0\alpha_3}^\uparrow \lambda_{0\beta_3}^\downarrow \lambda_{0\alpha_4}^\uparrow \lambda_{0\beta_4}^\downarrow. \end{aligned} \quad (3.21)$$

Moreover, we can construct nonzero-energy states containing even number of spinors:

- a state with the energy $-\mu$,

$$\lambda_{0\alpha}^\uparrow \lambda_{0\beta}^\uparrow \lambda_{0\gamma}^\uparrow \lambda_{0\delta}^\uparrow, \quad (3.22)$$

- $6 + 16 + 6 = 28$ states with the energy $-\mu/2$,

$$\lambda_{0\alpha}^\uparrow \lambda_{0\beta}^\uparrow, \quad \lambda_{0\alpha}^\uparrow \lambda_{0\beta}^\uparrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow, \quad \lambda_{0\alpha}^\uparrow \lambda_{0\beta}^\uparrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow, \quad (3.23)$$

- $6 + 16 + 6 = 28$ states with the energy $+\mu/2$,

$$\lambda_{0\alpha}^\downarrow \lambda_{0\beta}^\downarrow, \quad \lambda_{0\alpha}^\downarrow \lambda_{0\beta}^\downarrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow, \quad \lambda_{0\alpha}^\downarrow \lambda_{0\beta}^\downarrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow, \quad (3.24)$$

- a state with the energy $+\mu$,

$$\lambda_{0\alpha}^\downarrow \lambda_{0\beta}^\downarrow \lambda_{0\gamma}^\downarrow \lambda_{0\delta}^\downarrow. \quad (3.25)$$

We can also construct the fermionic sector of the supergravity multiplet as follows:

- $4 + 4 = 8$ fermionic states with the energy $-3\mu/4$,

$$\lambda_{0\alpha}^\uparrow \lambda_{0\beta}^\uparrow \lambda_{0\gamma}^\uparrow, \quad \lambda_{0\alpha}^\uparrow \lambda_{0\beta}^\uparrow \lambda_{0\gamma}^\uparrow \lambda_{0\alpha_1}^\downarrow \lambda_{0\beta_2}^\downarrow, \quad (3.26)$$

- $4 + 24 + 24 + 4 = 56$ fermionic states with the energy $-\mu/4$,

$$\lambda_{0\alpha}^\uparrow, \quad \lambda_{0\alpha}^\uparrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow, \quad \lambda_{0\alpha}^\uparrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow, \quad \lambda_{0\alpha}^\uparrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow \lambda_{0\alpha_3}^\uparrow \lambda_{0\beta_3}^\downarrow, \quad (3.27)$$

- $4 + 24 + 24 + 4 = 56$ fermionic states with the energy $+\mu/4$,

$$\lambda_{0\alpha}^\downarrow, \quad \lambda_{0\alpha}^\downarrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow, \quad \lambda_{0\alpha}^\downarrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow, \quad \lambda_{0\alpha}^\downarrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_1}^\downarrow \lambda_{0\alpha_2}^\uparrow \lambda_{0\beta_2}^\downarrow \lambda_{0\alpha_3}^\uparrow \lambda_{0\beta_3}^\downarrow, \quad (3.28)$$

- $4 + 4 = 8$ fermionic states with the energy $+3\mu/4$,

$$\lambda_{0\alpha}^\downarrow \lambda_{0\beta}^\downarrow \lambda_{0\gamma}^\downarrow, \quad \lambda_{0\alpha}^\downarrow \lambda_{0\beta}^\downarrow \lambda_{0\gamma}^\downarrow \lambda_{0\alpha_1}^\uparrow \lambda_{0\beta_2}^\downarrow. \quad (3.29)$$

The resulting spectrum is listed in Table. 1. In the pp-wave case, these states are splitting with an energy difference $\mu/4$. The resulting $2(1 + 28 + 35) = 128$ bosonic and the $2(8 + 56) = 128$ fermionic states correspond to the bosonic and the fermionic sectors of the supergravity multiplet in the flat case. It is confirmed that the above bosonic and the fermionic states

Energy	$-\mu$	$-\frac{3}{4}\mu$	$-\frac{1}{2}\mu$	$-\frac{1}{4}\mu$	0	$+\frac{1}{4}\mu$	$+\frac{1}{2}\mu$	$+\frac{3}{4}\mu$	$+\mu$
Boson	1		28		35 + 35		28		1
Fermion		8		56		56		8	

Table 1: Spectrum of the fermionic zero-mode Hamiltonian: The numbers of the states at each energy level are listed. The above states form a supermultiplet under the action of the kinematical supersymmetries.

correspond to the supergravity multiplet in the flat case by noting that such states appear in the expansion of the ground-state wave function in the flat case as $\lambda_\alpha \lambda_\beta \dots = (\lambda_\alpha^\dagger + \lambda_\alpha^\downarrow)(\lambda_\beta^\dagger + \lambda_\beta^\downarrow) \dots$. These are the coefficients in front of the plane-wave in the flat case. If we take the $\mu \rightarrow 0$ limit, then the above states with non-zero energies should have zero energies, that is, all of them should be degenerate. As a result, a $128 + 128$ supergravity multiplet at the rest frame is recovered.

Also, the resulting spectrum has the symmetry under $\mu \rightarrow -\mu$ as noted in Ref. [8]. Moreover, the spectrum resembles that of superstring theories. That is, $1 + 28 + 35$ or $8 + 56$ looks like the spectrum of NS-NS, R-R or R-NS sector, although we cannot give any definite physical interpretation.

Furthermore, the fermionic Hamiltonian is not positive definite, but it is bounded as

$$-\mu \leq H_F^{(0)} \leq +\mu. \quad (3.30)$$

The bosonic zero-mode Hamiltonian has a lower bound as denoted in Eq. (3.16) due to the presence of the zero-point energy $|\mu|$. Thus, we can easily find that the total zero-mode Hamiltonian is positive definite

$$H^{(0)} \geq 0.$$

Thus, the zero-energy ground state of the total zero-mode Hamiltonian can be uniquely constructed as[‡]

$$\begin{aligned}
\Psi_0^{(0)} &= \Psi_{B,\mu}^{(0)} \otimes \Psi_{F,-\mu}^{(0)} \\
&= \left(\frac{\mu}{3\pi}\right)^{3/4} \left(\frac{\mu}{6\pi}\right)^{3/2} \lambda_{0\alpha}^\dagger \lambda_{0\beta}^\dagger \lambda_{0\gamma}^\dagger \lambda_{0\delta}^\dagger \\
&\quad \times \exp \left(-\frac{1}{2} \cdot \frac{\mu}{3} \sum_{I=1}^3 (X_0^I)^2 - \frac{1}{2} \cdot \frac{\mu}{6} \sum_{m=4}^7 (X_0^m)^2 - \frac{\mu}{6} |Z_0|^2 \right), \quad (3.31)
\end{aligned}$$

[‡]We have fixed μ positive. If μ is negative, then $\lambda^\dagger$'s are replaced by $\lambda^\downarrow$'s.

which satisfies

$$H^{(0)}\Psi_0^{(0)} = 0.$$

Also, we can directly see that the above ground-state wave function is supersymmetric, i.e., $Q_\alpha^{(0)}\Psi_0^{(0)} = Q_\alpha^{(0)\dagger}\Psi_0^{(0)} = 0$ when the zero-modes of the dynamical supercharges are rewritten in terms of annihilation and creation operators as follows:

$$Q_\alpha^{(0)} = \left(e^{\frac{\mu}{12}\Gamma_{123}\tau}\right)_{\alpha\beta} \left[i\sqrt{2} \left(\sqrt{\frac{\mu}{3}} a_0^{I\dagger}(\Gamma_I)_{\beta\gamma} + \sqrt{\frac{\mu}{6}} a_0^{m\dagger}(\Gamma_m)_{\beta\gamma} \right) \lambda_{0\gamma}^\dagger \right. \\ \left. - i\sqrt{2} \left(\sqrt{\frac{\mu}{3}} a_0^I(\Gamma_I)_{\beta\gamma} + \sqrt{\frac{\mu}{6}} a_0^m(\Gamma_m)_{\beta\gamma} \right) \lambda_{0\gamma}^\dagger - 2\sqrt{\frac{\mu}{6}} \left(b_0 \frac{\partial}{\partial \lambda_{0\gamma}^\dagger} - b_0^\dagger \frac{\partial}{\partial \lambda_{0\gamma}^\dagger} \right) \right], \quad (3.32)$$

$$Q_\alpha^{(0)\dagger} = \left(e^{-\frac{\mu}{12}\Gamma_{123}\tau}\right)_{\alpha\beta} \left[i\sqrt{2} \left(\sqrt{\frac{\mu}{3}} a_0^I(\Gamma_I)_{\beta\gamma} + \sqrt{\frac{\mu}{6}} a_0^m(\Gamma_m)_{\beta\gamma} \right) \lambda_{0\gamma}^\dagger \right. \\ \left. - i\sqrt{2} \left(\sqrt{\frac{\mu}{3}} a_0^{I\dagger}(\Gamma_I)_{\beta\gamma} + \sqrt{\frac{\mu}{6}} a_0^{m\dagger}(\Gamma_m)_{\beta\gamma} \right) \lambda_{0\gamma}^\dagger - 2\sqrt{\frac{\mu}{6}} \left(b_0^\dagger \frac{\partial}{\partial \lambda_{0\gamma}^\dagger} - b_0 \frac{\partial}{\partial \lambda_{0\gamma}^\dagger} \right) \right]. \quad (3.33)$$

One can see that the ground-state wave function (3.31) is a singlet under the $SO(3) \times SO(6)$ as shown in Appendix A.

Now, we shall recall that the algebra of kinematical supersymmetries can be represented by[§]

$$Q^- = \lambda_0, \quad Q^{-\dagger} = \frac{\partial}{\partial \lambda_0}. \quad (3.34)$$

We can see that the above bosonic and the fermionic states form a supermultiplet under the action of the kinematical supersymmetries. When we decompose the representation of the kinematical supercharges as

$$Q^- = Q^{-\uparrow} + Q^{-\downarrow}, \\ Q^{-\dagger} = (Q^{-\uparrow})^\dagger + (Q^{-\downarrow})^\dagger,$$

$Q^{-\uparrow}$ and $(Q^{-\downarrow})^\dagger$ ($Q^{-\downarrow}$ and $(Q^{-\uparrow})^\dagger$) lower (raise) the fermionic energy. It should be remarked that the states with different energies belong to a multiplet because the supercharges do not commute with the Hamiltonian.

The story does not end here and we have to discuss the contribution of nonzero-modes (i.e., off-diagonal elements of the matrix variables), which will be studied in the next section.

[§]The factor i can be absorbed into the infinitesimal parameter of the kinematical supersymmetry transformation.

4 Nonzero-mode Ground-state Wave Function

In this section, we discuss the nonzero-mode Hamiltonian, its ground-state wave function and evaluate the L^2 -norm of the ground state.

To begin, we shall consider the abelian part of the nonzero-mode Hamiltonian. According to the similar discussion to that on the zero modes, we can easily show the following bounds:

$$H_B^C \geq |\mu|(N-1), \quad -|\mu|(N-1) \leq H_F^C \leq +|\mu|(N-1), \quad (4.1)$$

where the lowest state of H_F^C is represented by

$$\prod_{A=1}^{N-1} \lambda_{A\alpha}^\dagger \lambda_{A\beta}^\dagger \lambda_{A\gamma}^\dagger \lambda_{A\delta}^\dagger.$$

Recalling that matrix variables are restricted to the Cartan subalgebra in the IR region, we can say that the nonzero-mode Hamiltonian is positive definite in the same way as the zero-mode Hamiltonian. Also, the associated ground-state wave function in the IR region is normalizable.

We would like to consider the ground-state wave function of the nonzero-mode Hamiltonian including the interactions of non-diagonal elements of matrix variables. For this purpose, we have to solve two differential equations $Q\Psi = Q^\dagger\Psi = 0$, but it is quite difficult to solve them fully. In particular, the bosonic degrees of freedom for the $SO(6)$ part seem difficult to study. So let us truncate the bosonic variables X^i and Z into X^I ($I = 1, 2, 3$) ($SO(3)$ part) and also the 4-component spinors $\lambda^\uparrow$ and $\lambda^\downarrow$ into the 2-component ones, respectively. The truncation of the fermionic degrees is needed for the cancellation of the zero-point energy in the abelian parts of the nonzero-mode Hamiltonian. Such truncations modify the bounds of the abelian parts in bosonic and in fermionic nonzero-mode Hamiltonians respectively as

$$H_B^C \geq \frac{1}{2}|\mu|(N-1), \quad -\frac{1}{2}|\mu|(N-1) \leq H_F^C \leq +\frac{1}{2}|\mu|(N-1), \quad (4.2)$$

and the lowest state of the fermionic zero-mode Hamiltonian becomes

$$\prod_{n=1}^{N-1} \lambda_{A\alpha}^\uparrow \lambda_{A\beta}^\downarrow.$$

Therefore the above truncation preserves the positive definiteness in the IR region.

Now, the associated differential equations become simple forms

$$\begin{aligned}
Q_\alpha \Psi &= \left(e^{\frac{\mu}{12} \Gamma_{123} \tau} \right)_{\alpha\beta} \left[\left(\frac{\partial}{\partial X_A^I} - \frac{\mu}{3} X_A^I - \frac{1}{2} f_{ABC} \epsilon_{IJK} X_B^J X_C^K \right) \cdot (-i(\Gamma_I)_{\beta\gamma} \lambda_{A\gamma}^\dagger) \right. \\
&\quad \left. + \left(\frac{\partial}{\partial X_A^I} + \frac{\mu}{3} X_A^I + \frac{1}{2} f_{ABC} \epsilon_{IJK} X_B^J X_C^K \right) \cdot (-i(\Gamma_I)_{\beta\gamma} \lambda_{A\gamma}^\dagger) \right] \Psi \\
&= 0.
\end{aligned} \tag{4.3}$$

and

$$\begin{aligned}
Q_\alpha^\dagger \Psi &= - \left(e^{-\frac{\mu}{12} \Gamma_{123} \tau} \right)_{\alpha\beta} \left[\left(\frac{\partial}{\partial X_A^I} + \frac{\mu}{3} X_A^I + \frac{1}{2} f_{ABC} \epsilon_{IJK} X_B^J X_C^K \right) \cdot \left(-i(\Gamma_I)_{\beta\gamma} \frac{\partial}{\partial \lambda_{A\gamma}^\dagger} \right) \right. \\
&\quad \left. + \left(\frac{\partial}{\partial X_A^I} - \frac{\mu}{3} X_A^I - \frac{1}{2} f_{ABC} \epsilon_{IJK} X_B^J X_C^K \right) \cdot \left(-i(\Gamma_I)_{\beta\gamma} \frac{\partial}{\partial \lambda_{A\gamma}^\dagger} \right) \right] \Psi \\
&= 0.
\end{aligned} \tag{4.4}$$

We can solve the differential equations (4.3) and (4.4), and obtain a special solution for the wave function of nonzero-modes

$$\Psi_G = \prod_{A=1}^{N^2-1} \lambda_{A\alpha}^\dagger \lambda_{A\beta}^\dagger \Psi_B, \tag{4.5}$$

$$\begin{aligned}
\Psi_B &= \left(\frac{\mu}{3\pi} \right)^{3(N^2-1)/4} \exp \left[-\frac{1}{2} \cdot \frac{\mu}{3} \sum_{I=1}^3 \sum_{A=1}^{N^2-1} (X_A^I)^2 \right. \\
&\quad \left. + \frac{1}{3!} \sum_{I,J,K=1}^3 \sum_{A,B,C=1}^{N^2-1} f_{ABC} \epsilon_{IJK} X_A^I X_B^J X_C^K \right].
\end{aligned} \tag{4.6}$$

This solution is supersymmetric and has zero energy. Moreover, we can easily see that this solution is an $SO(3) \times SO(6)$ singlet and satisfies the constraint condition (2.59). However the function Ψ_B would not be square-integrable. In order to discuss the normalizability of this solution, let us set $N = 2$ (i.e., $SU(2)$ matrix model) for simplicity and evaluate the L^2 -norm of the above ground-state wave function $||\Psi_G||$ as

$$\begin{aligned}
||\Psi_G||^2 &\equiv \int \prod_{I,A} dX_A^I \Psi_G^\dagger \Psi_G \\
&= \int_{-\infty}^{\infty} dX_1^1 \int_{-\infty}^{\infty} dX_1^2 \cdots \int_{-\infty}^{\infty} dX_3^3 |\Psi_B|^2 \\
&= \left(\frac{\mu}{3\pi} \right)^{9/2} \times I,
\end{aligned} \tag{4.7}$$

where the I is defined by

$$\begin{aligned}
I &= \int_{-\infty}^{\infty} dX_1^1 \int_{-\infty}^{\infty} dX_1^2 \cdots \int_{-\infty}^{\infty} dX_3^3 e^{2f_0} e^{2f_1}, \\
f_0 &\equiv -\frac{1}{2} \cdot \frac{\mu}{3} \sum_{A,I=1}^3 (X_A^I)^2 \\
&= -\frac{1}{2} \cdot \frac{\mu}{3} \left[(X_1^1)^2 + (X_1^2)^2 + (X_1^3)^2 + (X_2^1)^2 \right. \\
&\quad \left. + (X_2^2)^2 + (X_2^3)^2 + (X_3^1)^2 + (X_3^2)^2 + (X_3^3)^2 \right], \\
f_1 &\equiv \frac{1}{3!} \sum_{A,B,C,I,J,K=1}^3 \epsilon_{ABC} \epsilon_{IJK} X_A^I X_B^J X_C^K \\
&= -X_1^3 X_2^2 X_3^1 + X_1^2 X_2^3 X_3^1 + X_1^3 X_2^1 X_3^2 \\
&\quad - X_1^1 X_2^3 X_3^2 - X_1^2 X_2^1 X_3^3 + X_1^1 X_2^2 X_3^3.
\end{aligned} \tag{4.8}$$

Here, we shall expand the f_1 part formally and evaluate the integral as follows:

$$I = \left(\frac{3\pi}{\mu} \right)^{9/2} \sum_{n=0}^{\infty} \frac{(2n+1)!}{n!} \left[\frac{1}{4} \left(\frac{3}{\mu} \right)^3 \right]^n. \tag{4.9}$$

This formula is obtained by calculating several terms with lower orders in the expansion by using the Mathematica. Though we have no analytical proof, we guess this formula (4.9) holds for general terms. As a result, we can evaluate the square integral of the bosonic ground-state as

$$||\Psi_G||^2 = 1 + O(1/\mu^3). \tag{4.10}$$

That is, the normalizability condition can be at least expressed as the asymptotic series which can be approximated with any amount of accuracy in the large μ limit. In fact, the convergence radius of this series is zero and in this sense we can say that this series is an asymptotic series. The n th partial sum of this asymptotic series does not converge as $n \rightarrow \infty$. However when μ is sufficiently large, we can take an n th partial sum for an appropriate finite n and calculate an approximate value of I . In this sense, we can write I as

$$I \sim 4\pi^{9/2} e^{-\left(\frac{\mu}{3}\right)^3} \int_0^{\left(\frac{\mu}{3}\right)^{3/2}} dt e^{t^2} - 2\pi^{9/2} \left(\frac{\mu}{3}\right)^{-3/2}.$$

Thus the corresponding wave function is not normalizable in the usual sense.

We note that the $n = 0$ part corresponds to the kinematical contribution and that $n \neq 0$ parts to the dynamical contributions arising from non-diagonal elements of matrix variables. In the IR region, the matrix variables are restricted to the Cartan subalgebra. Therefore the system can be described by the abelian part of the Hamiltonian and the ground state is square-integrable. If we consider a situation that the system flows from the IR region to the UV one, then it might be understood that the dynamical effects coming from non-diagonal components of matrix variables are turned on and tend to spoil down the normalizability of the ground-state wave function. If we take the limit $\mu \rightarrow \infty$, then we could sufficiently neglect contributions of off-diagonal elements (dynamical effects). It is also noted that the above series tends to diverge in the flat limit $\mu \rightarrow 0$.

Finally, we have a comment toward the full ground-state wave function. In the above derivation, we have truncated the bosonic and fermionic variables. But, it would be possible to avoid the truncation of the fermionic degrees. In this case, it seems that the numbers of the bosons and fermions are asymmetric. Our interpretations are as follows: First, it might be possible to decompose the ground-state wave function into the bosonic and fermionic parts as $\Psi = \Psi_B \otimes \Psi_F$. It is no wonder since the fermionic Hamiltonian contains mass term and the well-known argument in the flat space would not hold on the pp-wave. Therefore, if we suppose that such decomposition is possible, then it is not necessary that the bosonic wave function is identically zero. Second, the fermionic ground state might be completely determined by the mass term. When we express the fermionic Hamiltonian by $H_F = H_F^{\text{Yukawa}} + H_F^{\text{mass}}$, the vacuum expectation value (VEV) is written as

$$(\Psi, H_F \Psi) = (\Psi, H_F^{\text{Yukawa}} \Psi) + (\Psi_F, H_F^{\text{mass}} \Psi_F). \quad (4.11)$$

The VEV of the Yukawa coupling term contains the term $(\Psi_B, X \Psi_B)$ and hence must vanish due to the requirement of an $SO(3) \times SO(6)$ symmetry for the ground state. On the other hand, it seems quite difficult to treat the full bosonic part. In particular, the treatment of the $SO(6)$ part seems hard to analyze. We also guess that the solution without the truncation of the fermionic variables could be obtained by setting X^m, Z to zero in the full solution.

5 Conclusions and Discussions

We have considered the ground-state wave function of the supermembrane on the pp-wave by introducing the complex-spinor notation ($Spin(7) \times U(1)$ formalism). The superalgebra of the

complex supercharges has been also calculated.

The zero-mode Hamiltonian and its ground state have been intensively studied. In particular, we have shown that the zero-mode Hamiltonian is positive definite and explicitly constructed the unique supersymmetric ground-state wave function as the product of ground states of bosonic harmonic oscillators and fermionic variables. This wave function is square-integrable. The ground state for the zero-mode is splitting in contrast to that of the flat case, and the zero-mode Hamiltonian has the string-like spectrum.

We have also investigated the nonzero-mode Hamiltonian and constructed an example of the supersymmetric ground-state wave function for the nonzero-mode with the truncation of variables in the $N = 2$ case. This ground state seems non-normalizable, but its L^2 -norm can be defined by an asymptotic series. This series diverges since its convergence radius is zero. The dynamical effects coming from non-diagonal elements of matrix variables are represented as corrections with order $1/\mu^3$. In the IR region, matrix variables are restricted to the abelian part, whose ground-state wave function is normalizable. We might expect that the interactions become stronger as the system flows from the IR region to the UV one. It might be considered that these dynamical contributions tend to spoil down the normalizability of the ground state, at least in our example of the ground state. It would be possible to obtain the full solution without the truncation. It would be nice to proceed the study of the nonzero-mode further. In particular, it is an interesting future problem to obtain the full nonzero-mode ground-state wave function. If this is possible, then one can diagonalize the nonzero-mode Hamiltonian and obtain the spectrum. Such results shed light upon the spectrum of the supermembrane in the flat space. Also, the extension of our example to an arbitrary N is an interesting problem.

There exist other interesting future problems. The study of the Witten index for the ground state [25] seems very interesting since the energy of bosons is different from that of the corresponding fermions because the supercharges do not commute with the Hamiltonian. It would be also nice to consider the supermembrane on less supersymmetric backgrounds as discussed in [26].

Acknowledgement

The work of K.S. is supported in part by the Grant-in-Aid from the Ministry of Education, Science, Sports and Culture of Japan (# 14740115).

Appendix

A Proof for the $SO(3) \times SO(6)$ Invariance of the Ground State

We shall prove that the ground-state wave function (3.31) of the zero-mode Hamiltonian is invariant under the $SO(3) \times SO(6)$ transformation. The orbital part of the total angular momenta trivially vanish due to the expressions (2.54)-(2.58). The remaining problem is to show that the spin part vanishes.

The spin part of (2.54) and (2.55) can be rewritten as

$$S^{IJ} = -\frac{i}{2}\lambda_0^\uparrow \Gamma^{IJ} \frac{\partial}{\partial \lambda_0^\uparrow} - \frac{i}{2}\lambda_0^\downarrow \Gamma^{IJ} \frac{\partial}{\partial \lambda_0^\downarrow}, \quad S^{mn} = -\frac{i}{2}\lambda_0^\uparrow \Gamma^{mn} \frac{\partial}{\partial \lambda_0^\uparrow} - \frac{i}{2}\lambda_0^\downarrow \Gamma^{mn} \frac{\partial}{\partial \lambda_0^\downarrow}, \quad (\text{A.1})$$

since the matrices Γ^{IJ} and Γ^{mn} commute with $i\Gamma_{123}$ and the action of these matrices preserves the chirality. The action of parts including $\lambda^\downarrow$ trivially annihilate the state. The spin parts including $\lambda^\uparrow$ replace the $\lambda_{0\alpha_0}$ with $\lambda_\alpha^\uparrow (\Gamma^{IJ})_{\alpha\alpha_0}$, $(\lambda_\alpha^\uparrow (\Gamma^{mn})_{\alpha\alpha_0})$. It should be noted that the matrices Γ^{IJ} or Γ^{mn} should map $\lambda^\uparrow$ to other $\lambda^\uparrow$ or zero since the Γ^{IJ} and Γ^{mn} are antisymmetric. Thus, the spin part leads to zero due to the overlap of $\lambda^\uparrow$'s.

Next, we shall consider the spin parts (2.56) and (2.57), which are rewritten as

$$S_0^{m+} = -\frac{1}{2}\lambda_0^\uparrow \Gamma^m \lambda_0^\downarrow, \quad S_0^{m-} = -\frac{1}{2} \frac{\partial}{\partial \lambda_0^\uparrow} \Gamma^m \frac{\partial}{\partial \lambda_0^\downarrow}. \quad (\text{A.2})$$

We can easily see that these spins vanish on the ground state (3.31).

Finally, let us consider the spin part of (2.58), which is given by

$$S_0^{89} = -\frac{1}{2}\lambda_0^\uparrow \frac{\partial}{\partial \lambda_0^\uparrow} - \frac{1}{2}\lambda_0^\downarrow \frac{\partial}{\partial \lambda_0^\downarrow} + 2. \quad (\text{A.3})$$

The part containing $\lambda^\downarrow$ trivially annihilates the state. The action of the part written by $\lambda^\uparrow$ leads to the eigenvalue -2 , which is canceled with the normal-ordering constant $+2$.

The extension of the above proof to the nonzero-mode ground-state wave function discussed in the manuscript is a simple exercise.

References

- [1] J. Kowalski-Glikman, “Vacuum states in supersymmetric Kaluza-Klein theory”, *Phys. Lett. B* **134** (1984) 194.
- [2] J. Figueroa-O’Farrill and G. Papadopoulos, “Homogeneous fluxes, branes and a maximally supersymmetric solution of M-theory”, *JHEP* **0108** (2001) 036, [hep-th/0105308](#).
- [3] R. Penrose, “Any spacetime has a plane wave as a limit”, *Differential geometry and relativity*, Reidel, Dordrecht, 1976, pp. 271-275.
R. Güven, “Plane Wave Limits and T-duality”, *Phys. Lett. B* **482** (2000) 255, [hep-th/0005061](#).
- [4] M. Blau, J. Figueroa-O’Farrill, C. Hall and G. Papadopoulos, “A new maximally supersymmetric background of IIB superstring theory”, *JHEP* **0201** (2001) 047, [hep-th/0110242](#).
“Penrose limits and maximal supersymmetry”, [hep-th/0201081](#).
- [5] R.R. Metsaev, “Type IIB Green-Schwarz superstring in plane wave Ramond-Ramond background”, *Nucl. Phys. B* **625** (2002) 70, [hep-th/0112044](#).
- [6] R.R. Metsaev and A.A. Tseytlin, “Exactly solvable model of superstring in plane wave Ramond-Ramond background ”, *Phys. Rev. D* **65** (2002) 126004, [hep-th/0202109](#).
- [7] J.G. Russo and A.A. Tseytlin, “On solvable models of type IIB superstring in NS-NS and R-R plane wave backgrounds ”, *JHEP* **0204** (2002) 021, [hep-th/0202179](#).
- [8] D. Berenstein, J. Maldacena and H. Nastase, “Strings in flat space and pp waves from $\mathcal{N} = 4$ Super Yang Mills”, *JHEP* **0204** (2002) 013, [hep-th/0202021](#).
- [9] E. Bergshoeff, E. Sezgin and P. Townsend, “Supermembranes and Eleven-Dimensional Supergravity”, *Phys. Lett.* **189** (1987) 75.
- [10] E. Bergshoeff, E. Sezgin and P. Townsend, “Properties of the Eleven-Dimensional Supermembrane Theory”, *Ann. Phys.* **185** (1988) 330.
- [11] B. de Wit, J. Hoppe and H. Nicolai, “On the Quantum Mechanics of Supermembranes”, *Nucl. Phys. B* **305** (1988) 545.

- [12] T. Banks, W. Fischler, S.H. Shenker and L. Susskind, “M-theory As A Matrix Model: Conjecture”, Phys. Rev. **D 55** (1997) 5112, [hep-th/9610043](#).
- [13] K. Dasgupta, M.M. Sheikh-Jabbari and M. Van Raamsdonk, “Matrix Perturbation Theory For M-Theory On a PP-Wave”, JHEP **0205** (2002) 056, [hep-th/0205185](#).
- [14] K. Sugiyama and K. Yoshida, “Supermembrane on the PP-wave Background”, to appear in Nucl. Phys. **B**, [hep-th/0206070](#).
- [15] T. Banks, N. Seiberg and S. Shenker, “Branes from Matrices”, Nucl. Phys. **B 490** (1997) 91, [hep-th/9612157](#).
- [16] K. Sugiyama and K. Yoshida, “BPS Conditions of Supermembrane on the PP-wave”, to appear in Phys. Lett. **B**, [hep-th/0206132](#).
- [17] N. Kim and J. Plefka, “On the Spectrum of PP-Wave Matrix Theory”, [hep-th/0207034](#).
K. Dasgupta, M.M. Sheikh-Jabbari and M. Van Raamsdonk, “Protected multiplets of M-theory on a Plane Wave”, [hep-th/0207050](#).
N. Kim and J.-H. Park, “Superalgebra for M-theory on a pp-wave”, [hep-th/0207061](#).
J.-H. Park, “Supersymmetric objects in the M-theory on a pp-wave”, [hep-th/0208161](#).
- [18] R.C. Myers, “Dielectric-Branes”, JHEP **9912** (1999) 022, [hep-th/9910053](#).
- [19] K. Sugiyama and K. Yoshida, “Giant Graviton and Quantum Stability in Matrix Model on PP-wave Background”, to appear in Phys. Rev. **D**, [hep-th/0207190](#).
- [20] D. Bak, “Supersymmetric Branes in PP Wave Background”, [hep-th/0204033](#).
S. Hyun and H. Shin, “Branes from Matrix Theory in PP-Wave Background”, to appear in Phys. Lett. **B**, [hep-th/0206090](#).
M. Alishahiha and M. Ghasemkhani, “Orbiting Membranes in M-theory on $AdS_7 \times S^4$ Background”, [hep-th/0206237](#).
N. Kim, K.M. Lee and P. Yi, “Deformed Matrix Theories with N=8 and Fivebranes in the PP Wave Background”, [hep-th/0207264](#).
A. Mikhailov, “Nonspherical Giant Gravitons and Matrix Theory”, [hep-th/0208077](#).
- [21] K. Sugiyama and K. Yoshida, “Type IIA String and Matrix String on PP-wave”, to appear in Nucl. Phys. **B**, [hep-th/0208029](#).

- [22] M.J. Duff, P.S. Howe, T. Inami and K.S. Stelle, “Superstrings in $D = 10$ from Supermembranes in $D = 11$ ”, Phys. Lett. **B 191** (1987) 70.
- [23] S. Hyun and H. Shin, “N=(4,4) Type IIA String Theory on PP-Wave Background”, hep-th/0208074.
- [24] B. de Wit, M. Lüscher and H. Nicolai, “The Supermembrane is Unstable”, Nucl. Phys. **B 320** (1989) 135.
- [25] P. Yi, “Witten Index and Threshold Bound States of D-Branes”, Nucl. Phys. **B505** (1997) 307, hep-th/9704098.
S. Sethi and M. Stern, “D-Brane Bound States Redux”, Commun. Math. Phys. **194** (1998) 675, hep-th/9705046.
- [26] K.M. Lee, “M-theory on Less Supersymmetric PP-Waves”. hep-th/0209009.